



Perspective paper

Ethnobotany/ethnopharmacology and mass bioprospecting: Issues on intellectual property and benefit-sharing

D.D. Soejarto^{a,b,*}, H.H.S. Fong^a, G.T. Tan^a, H.J. Zhang^a, C.Y. Ma^a, S.G. Franzblau^a, C. Gyllenhaal^a, M.C. Riley^a, M.R. Kadushin^{a,b}, J.M. Pezzuto^c, L.T. Xuan^d, N.T. Hiep^d, N.V. Hung^d, B.M. Vu^d, P.K. Loc^d, L.X. Dac^d, L.T. Binh^d, N.Q. Chien^d, N.V. Hai^d, T.Q. Bich^e, N.M. Cuong^e, B. Southavong^f, K. Sydara^f, S. Bouamanivong^f, H.M. Ly^g, Tran Van Thuy^h, W.C. Roseⁱ, G.R. Dietzman^j

^a PCRPS, College of Pharmacy, University of Illinois at Chicago, 833 S. Wood Street, Chicago, IL 60612, USA

^b Field Museum, Chicago, IL, USA

^c Schools of Pharmacy, Nursing, and Health Sciences, Purdue University, West Lafayette, IN, USA

^d Vietnamese Academy of Science and Technology, Hanoi, Vietnam

^e Cuc Phuong National Park, Ninh Binh, Vietnam

^f Traditional Medicine Research Center, Vientiane, Laos

^g Laboratory of Mycobacteria, National Institute of Hygiene and Epidemiology, Hanoi, Vietnam

^h Department of Botany, Hanoi University of Science, Hanoi, Vietnam

ⁱ Bristol-Myers Squibb Pharmaceutical Research Institute, Princeton, NJ, USA

^j White Point Systems, Friday Harbor, WA, USA

Accepted 23 May 2005

Abstract

Ethnobotany/ethnopharmacology has contributed to the discovery of many important plant-derived drugs. Field explorations to seek and document indigenous/traditional medical knowledge (IMK/TMK), and/or the biodiversity with which the IMK/TMK is attached, and its conversion into a commercialized product is known as bioprospecting or biodiversity prospecting. When performed in a large-scale operation, the effort is referred to as mass bioprospecting. Experiences from the mass bioprospecting efforts undertaken by the United States National Cancer Institute, the National Cooperative Drug Discovery Groups (NCDDG) and the International Cooperative Biodiversity Groups (ICBG) programs demonstrate that mass bioprospecting is a complex process, involving expertise from diverse areas of human endeavors, but central to it is the Memorandum of Agreement (MOA) that recognizes issues on genetic access, prior informed consent, intellectual property and the sharing of benefits that may arise as a result of the effort. Future mass bioprospecting endeavors must take heed of the lessons learned from past and present experiences in the planning for a successful mass bioprospecting venture.

© 2005 Elsevier Ireland Ltd. All rights reserved.

Keywords: Ethnobotany; Ethnopharmacology; Mass bioprospecting; NCI; NCDDG; UIC ICBG; Intellectual property; Prior informed consent; Indigenous and traditional knowledge; Benefit-sharing

1. Introduction

The definition of the term ethnobotany, originally applied to the study of the utilitarian relationship [relationship that includes use for medicinal purposes] between humans and the plant environment in primitive settings (Harshberger, 1896),

* Corresponding author. Tel.: +1 312 996 8889; fax: +1 312 413 5894.

E-mail address: dds@uic.edu (D.D. Soejarto).

has now evolved into a much broader meaning that covers not only a utilitarian relationship, but also relationships that embrace the symbolic, ecological and cognitive, as well as the human–plant relationship in a modern setting (Schultes and von Reis, 1995; Alexiades, 1996). On the other hand, the more recent term ethnopharmacology has undergone only slight evolution in meaning since its original definition as “a multi-disciplinary area of research, concerned with the observation, description and experimental investigation of indigenous drugs and their biological activities” (Rivier and Bruhn, 1979). Its contemporary definition addresses the “interdisciplinary study of the physiological actions of plant, animal and other substances used in indigenous medicines of past and present cultures” (International Society of Ethnopharmacology Constitution, 2005). The breath of studies embraces “use of plants, fungi, animals, microorganisms and minerals and their biological and pharmacological effects based on the principles established through international convention”, as well as “the observation and experimental investigation of the biological activities of plant and animal substances”, using ethnopharmacological, ethnobotanical, ethnochemical, pharmacological and toxicological approaches (Journal of Ethnopharmacology, 2005).

Since these two branches of scientific disciplines carry a common denominator, namely the human or cultural component inherent in the word “ethno”, ethnobotanical and ethnopharmacological studies normally involve field explorations of indigenous as well as traditional medical knowledge (IMK/TMK), together with the biodiversity component to which such knowledge is attached, documentation and conversion of the knowledge into a product, be it a scholarly paper, a book, photographic images or a tangible product of commercial value. In this process, when the purpose and outcome are commercial in nature, the activity is known today as “bioprospecting”, a concept originally introduced in 1989 for “chemical prospecting” (Eisner, 1989, 1992) and was re-defined in 1993 as “biodiversity prospecting” (Reid et al., 1993). The term applies also to the exploration and utilization of the biological diversity itself for commercial purposes, either within the context of IMK/TMK or without.

Twenty-five years ago, when the *Journal of Ethnopharmacology* was established (*Journal of Ethnopharmacology* 1 (1) 1979), bioprospecting was considered natural and justified, because the outcome benefits the scientific and general communities, locally and globally. In fact, bioprospecting has been going on for centuries, but only within the past 20 years has this activity gained prominence due to technological advances in pharmaceutical, biotechnological and agricultural sectors. Before this period, little thought was given to the issue of ownership of IMK and TMK and of the distribution of benefits that may arise as the result of the bioprospecting effort. This had been due, in part, to the general attitude at that time that the world’s biodiversity, namely the genetic or biological resources with which IMK/TMK is attached, represents a common heritage of man (FAO, 1983) and so does the IMK/TMK.

2. Recognition of IMK/TMK and “sovereign rights” over genetic resources

The turning point on the formal recognition of ownership of indigenous and traditional knowledge took place in 1988 at the first International Congress of Ethnobiology (ICE) in Belém (Brazil), with the “Declaration of Belém” (<http://guallart.dac.uga.edu/ISE/iseBelem.html>), which is embodied within the Code of Ethics of the International Society of Ethnobiology (2005) (<http://guallart.dac.uga.edu/ISE/iseEthics.html>). The Code of Ethics states that “indigenous peoples, traditional societies and local communities have a right to self determination (or local determination for traditional and local communities) and that researchers and associated organizations will acknowledge and respect such rights” (Principle 2) and that they “must be fairly and adequately compensated for their contribution to ethnobiological research activities and outcomes involving their knowledge” (Principle 12). In the context of IMK/TMK, IMK/TMK represents the property of the holders of that knowledge and must be respected, and that compensation to the holders must be provided for the utilization and conversion of such knowledge (bioprospecting) into a tangible product. In the second ICE in 1991, in Kunming, PRC, participants went further to declare that traditional and indigenous knowledge are *inventive* and *intellectual*, therefore, worthy of protection in all legal, ethical and professional frameworks (International Society of Ethnobiology Constitution, 2005).

In 1992, the recognition of ownership of traditional knowledge and the biodiversity to which the knowledge is attached, and the sharing of the benefits that may arise as a result of their utilization was enforced in the form of the United Nations Convention on Biodiversity (Secretariat of the Convention on Biological Diversity, 2001, Articles 15.1 and 8.j).

The Belém Declaration, the Kunming Action Plan, the United Nations Convention on Biodiversity and numerous other Declarations of similar nature represent the instruments for States to legislate the utilization and protection of their natural resources that include the biological diversity. One of the earlier biodiversity policy regulations enacted by a country was the Executive Order 247 (EO 47) (1995) introduced by the President of the Republic of the Philippines. Other countries have also enacted legislation or are in the process of regulating their policy. Numerous national and international forums have been convened toward the legislation of a national bioprospecting policy. Central to any policy on bioprospecting are the issues on intellectual property, as it pertains to IMK/TMK, and the benefit-sharing of the process and the outcome that may arise from the activity.

Bioprospecting is performed by a diverse class of people, non-scientists and scientists alike. Although the context of this paper is ethnobotany and ethnopharmacology, the discussion that follows covers bioprospecting activities triggered by the search and utilization of IMK/TMK, as well as by the search and utilization of the component to which IMK/TMK is attached. When the activity of bioprospecting involves a

large-scale effort to search and commercialize IMK/TMK and/or the biological diversity, this effort is referred to as “mass bioprospecting”.

3. Mass bioprospecting: contemporary models

The effort of the United States National Cancer Institute (NCI) in searching for anticancer agents from the biodiversity, in the present context, of plants, is one example of mass bioprospecting. In this effort, field explorations are largely guided by the so-called biodiversity or “random” collection approach, with ethnobotanical or ethnopharmacological information playing a minimal or no role. NCI launched its effort in 1955, and for the period of 1960–1982, about 114,000 extracts from an estimated 35,000 plant samples (representing 12,000–13,000 species) collected mostly from temperate regions of the world had been screened against a number of tumor systems (Cragg and Boyd, 1996; Cragg et al., 1996). A wide variety of compound classes were isolated and characterized. Clinically significant cancer chemotherapeutic agents that emerged from this program included paclitaxel (*Taxus brevifolia* Nutt. and other *Taxus* species, Taxaceae), hycamptamine (topotecan), CPT-11 and 9-aminocamptothecin. The latter three compounds are semisynthetic derivatives of camptothecin (*Camptotheca acuminata* Decne., Nyssaceae) (Cragg et al., 1993). The program was extended from 1986 to 2004, with an emphasis on global plant collections and screening against tumor cell cultures. Although no provision was made in the recognition of intellectual property and in the arrangement of benefit-sharing during the first phase of the NCI's effort (1955–1980), a “Letter of Intent”, later evolving into a “Letter of Collection” (LOC), then a Memorandum of Agreement (MOA), was in place as an umbrella for the field operation of the NCI contractors in the second phase of NCI-sponsored explorations (1986–2004). The LOI/LOC/MOA contains provisions to recognize the ownership of genetic resources and an arrangement to share the benefits of discovery (Mays and Mazan, 1996; Hallock and Cragg, 2003).

In 1983, the NCI's mass bioprospecting effort was extended through the establishment of a National Cooperative Drug Discovery Group (NCDDG) program by the Developmental Therapeutics Program (DTP), Division of Cancer Treatment and Diagnosis (DCT). This program supports broad, innovative and multi-disciplinary approaches to the discovery of new, synthetic or natural-source-derived anticancer agents (Hallock and Cragg, 2003; NCI-DTP-DCTGCOB, 2005). In the programs that target biodiversity as the source of discovery, biodiversity-based collection approach also formed the basis of organism selection. In the NCDDG program, issues on IP and benefit-sharing were dealt with individually by each group. Recently, the overall effort and accomplishments of the NCDDG projects were reviewed (Hallock and Cragg, 2003).

Commencing in 1993, the International Cooperative Biodiversity Groups (ICBG), a program administered by the Fogarty International Center (FIC), National Institutes of Health (NIH) and supported through funds from NIH, National Science Foundation (NSF) and US Department of Agriculture (USDA) Foreign Agricultural Service (FAS), started operation in an effort to integrate the goals of improvement of human health through drug discovery, incentives for conservation of biodiversity and development of new models of sustainable economic activity that focus on the environment, health, equity and democracy. The implementation of this program is based on the belief that the discovery and development of pharmaceutical and other useful agents from the world's biodiversity can, under appropriate circumstances, promote scientific capacity development and economic incentives to conserve the biological resources from which these products are derived (Fogarty International Center, 2004). This unique effort is undertaken in such a way that local communities and other source country organizations can derive direct benefits from the effort and, ultimately, from their diverse biological resources, so that benefit-sharing may provide clear incentives for preservation and sustainable use of the biodiversity (Rosenthal, 1996, 1997; Rosenthal et al., 1999; Fogarty International Center, 2004).

The ICBG's drug discovery effort is targeted to a broad range of organisms, including five of the six recognized kingdoms of biodiversity: Eubacteria, Protoctista, Plantae, Fungi and Animalia (NIH News, 2003).

4. “Studies on biodiversity of Vietnam and Laos”: an ICBG program based at the University of Illinois at Chicago

A complete review of the ICBG has been published (Rosenthal and Pezzuto, 1999). The University of Illinois at Chicago (UIC)-based ICBG, known simply as “UIC ICBG” or “Studies on Biodiversity of Vietnam and Laos” (Soejarto et al., 1999, 2002a,b, 2004a,b), serves as a model for the implementation of the ICBG principles, which are, ultimately, the principles of the United Nations Convention on Biological Diversity. In its current Phase II operation (2003–2008), this ICBG consortium consists of two US-based academic institutions (UIC and Purdue University), two Vietnamese research institutions (Vietnamese Academy of Science and Technology; and Cuc Phuong National Park), one Lao research institution (Traditional Medicine Research Center) and an industrial partner (Bristol-Myers Squibb, BMS). Although the drug discovery and development objective of the UIC-ICBG is to uncover biologically active molecules from plants of Vietnam and Laos as chemotherapeutic candidates for malaria, tuberculosis, AIDS, cancer and CNS-related diseases, the bioprospecting goals are more all-encompassing, namely: (a) to discover and develop new drugs from plants of Vietnam and Laos (as just stated); (b) to promote economic development among communities in the ICBG study areas

and to promote institutional-infrastructure strengthening and human-resource development of host-country institutions; (c) to undertake biotic survey and to promote conservation initiatives in Vietnam and Laos. In other words, the UIC ICBG is a bioprospecting endeavor that is truly multi-disciplinary and multi-national, and truly integrates drug discovery, biodiversity conservation and economic development. Since space only allows a rather narrow discussion, addressing, specifically, the drug discovery and development activities, readers are directed to papers already published that document activities in response to the other two goals of the UIC ICBG's bioprospecting effort (Hiep and Loc, 2000; Riley, 2001; Soejarto et al., 2001, 2002c,d, 2004a,b; Dietzman et al., 2002; Hiep et al., 2002; Loc et al., 2002, 2004; Vu et al., 2002a,b; Bien et al., 2003, 2004; Dzu et al., 2003; TMRC, 2003; Dac et al., 2004; ICBG/AP4, 2005).

In drug discovery and development, approaches utilized in the selection of plants consist of biodiversity-based collection ["random" collection] centered in the Cuc Phuong National Park in Vietnam, and ethnobotany-driven interviews on the medicinal uses of plants in Laos. Ethnobotanical interviews were also performed in Vietnam during the first Phase of the UIC ICBG project (1998–2003), among communities (Muong ethnic minority) in villages surrounding CPNP.

Seven years after the start of the UIC ICBG project (1998–2005), 3331 plant samples have been collected from Vietnam (Cuc Phuong National Park/CPNP) based on biodiversity ("random") approach, comprising >950 species of flowering plants identified to species level. Since the results of the taxonomic inventory of plants in this park indicate that 1926 species of Angiosperms (Soejarto et al., 2004c) are found at CPNP, >900 species still remain to be collected and screened. In addition, 960 plant samples (about >700 species) have been collected based on ethnobotanical/ethnopharmacological field interviews with Muong communities surrounding CPNP in Vietnam, and with ethnic communities throughout Laos. For every plant sample collected, a set of seven voucher herbarium specimens was collected for deposit at the consortium's herbaria and at other herbaria of collaborating botanical institutions.

The drug discovery process in the UIC ICBG follows classic pharmacognosy methods, through sample extraction, high-throughput screening of extracts in the target disease system, recollection of samples (same plant parts) of active species and bioassay-guided fractionation and isolation. CH_2Cl_2 -soluble extracts or CHCl_3 - or CH_2Cl_2 -soluble fractions from MeOH/EtOH extracts are assayed, after elimination of tannins and polyphenols, which are known to interfere with certain enzyme-based assays.

Extracts that have shown confirmed activity are fractionated/isolated using an effective combination of expertise and advanced separation technology protocol, typically based on the combination of flash (Isco CombiFlash SG100C Separation System), semi-preparative and preparative HPLC chromatography, while droplet counter-current systems are employed to maximize efficiency. Traditional

methods include vacuum, gravity, flash or low–medium pressure column chromatography using a variety of adsorbents and media. Early LC/MS/MS-based dereplication strategies that favor novel bioactive molecules over non-specific, known and/or unwanted compounds from mixtures are employed to support the bioassay-directed fractionation and isolation process. Chemical profiling by HPLC–MS is used, especially, for chemotaxonomically related species under investigation.

State-of-the-art physical and spectroscopic methods, such as high field nuclear magnetic resonance spectrometry (NMR), time-of-flight mass spectrometry (TOF-MS), chemical ionization (CI), field desorption and fast-atom bombardment (FAB) MS, are employed in structure elucidation of active compounds. Stereochemistry is determined by the use of ORD and CD, coupled with high-field ^1H and ^{13}C NMR spectroscopy, with appropriate two-dimensional (2D) and decoupling experiments. The use of single crystal X-ray crystallographic analysis is routinely performed to solve difficult stereochemistry assignments.

At the end, new compounds that are routinely isolated based on a particular biological activity are included in the UIC ICBG natural products library and incorporated in the NAPIS Database, the computerized documentation system utilized in the ICBG program, for periodic submission to the ICBG's Global Data Center (GDC) based in Friday Harbor, Washington State. These compounds may eventually be exposed to multiple screens across the ICBGs.

5. UIC ICBG bioprospecting effort: results

Of the 3331 samples collected, extracts of 2309 (comprising approximately 800 species) have been screened in anti-HIV, antimalaria and anti-TB disease systems, as well as in a tumor cell lines panel. Twenty-two recollections of active samples have been made and approximately 280 pure natural products of varying degrees of structural complexity and/or biological activity have been isolated. Aside from the discovery of biologically active compounds, the UIC ICBG studies have also made significant contribution to the knowledge of natural products chemistry through the isolation and elucidation of 80 new secondary metabolites and novel chemical entities, reported for the first time from higher plants. This high number may have been the result of rigorous prioritization criteria in the project, based on pre-existing biological and chemical information (both at species and generic levels), as well as on a chemical dereplication strategy using the resources and expertise available at UIC. The chemical diversity of these new natural products includes alkaloids/amides, macrocyclics, lignans, neolignans, butenolides, phenylpropanoids, terpenes, norditerpenes, triterpenes and steroids (Zhang et al., 2001, 2002a,b; Chien et al., 2004).

Ten of the 80 new chemical entities have a novel carbon skeleton, being described and communicated for the first time, all of which are structurally related anti-HIV sesquiterpenes (litseaverticillols A–J) isolated from *Litsea verticillata*

Hance (Lauraceae). These compounds are unique, because they are α,β -conjugated pentacyclosesquiterpenes with a nine-member side chain (Zhang et al., 2003, in press). A number of the active anticancer compounds are in various stages of preclinical testing, including evaluation in animal models.

Central to the existence and operation of the UIC ICBG is the Memorandum of Agreements (1999–2005, 2004–2010), which spell out issues on access to genetic resources and to IMK/TMK, recognition of intellectual property rights on discovery and on IMK/TMK, and the sharing of benefits as part of the bioprospecting process, as well as of the benefits that may materialize (long-term benefits, namely royalties), as a result of the bioprospecting effort (Soejarto et al., 2004a,b).

6. Mass bioprospecting in the future

Estimates place the number of species of organisms at between 1,392,485 (Wilson, 1988) and 1,750,000 (Hammond, 1995), classified into Kingdoms Eubacteria (bacteria, cyanobacteria “blue-green algae”), Archaea (halobacterians, methanogens, eocytes), Protoctista (protozoa, “algae”), Plantae (land plants), Fungi (molds, lichen-forming, yeasts, mushrooms) and Animalia (mesozoa, invertebrates, mammals) (Hammond, 1995).

Examination of the statistics on number of described species shows that arthropods (Kingdom Animalia), comprising about 1,085,000 species, including insects, make up the largest portion (75.4%) of the earth's biodiversity (Hammond, 1995). Insects alone, estimated at 950,000 species, comprise 62% of global biodiversity. Despite the fact that the number of insect species is estimated to be between 2,000,000 and 100,000,000 (Hammond, 1995), this group remains relatively untouched as a source of novel compounds by the drug discovery community. Animal species [Kingdom Animalia], excluding the arthropods, comprise 13%, while the fungi and the protists [protoctists] represent 4% and 5% of global biodiversity, respectively. It is reasonable to assume that the overwhelming majority of fungal species have yet to be isolated and tested for the production of biologically active compounds, even though members of the order Actinomycetales, like Streptomycetes, have been the most prominently known microbial producers of natural products.

Based on the number of known species (300,000–500,000) (Hammond, 1995), plants represent the second largest source of biodiversity (15%). Estimates have been made that between 20,000 and 55,000 species of plants have been used medicinally (Penso, 1976; Schippmann et al., 2002), of which only a small portion has been investigated for drug purposes. Among those that have been investigated are plant species that produce important drugs such as quinine, reserpine, tubocurarine, vincristine, vinblastine, pilocarpine, atropine, morphine, cocaine, to mention a few. Overall, only 15–20% of terrestrial plants have been evaluated for pharmaceutical potential. Consequently, plants, including uninvestigated

medicinal plants, continue to represent a significant pool of raw material for the discovery of new drugs.

With the biodiversity statistics presented above, the world's biodiversity and the ethnobotanical and ethnopharmacological treasure house that remains in store will continue to present an attractive target for future mass bioprospecting effort.

7. Discussion and conclusions

From the experiences gained through the participation in the NCI (Soejarto, 1993; Soejarto et al., 1996), NCDDG (Kingham et al., 2003) and ICBG (Soejarto et al., 1999, 2002a,b, 2004a,b) bioprospecting projects described above, clearly, such endeavors are a highly complex process. Future mass bioprospecting effort must incorporate lessons learned from these experiences. The most important consideration is the broad spectrum of requirements that must be amalgamated: team scientific expertise (of all relevant disciplines) together with expertise in a wide range of human endeavors, including diplomacy, international laws and legal understandings, social sciences, politics, anthropology and good common sense. Equally important is the fact that such endeavors must be international in nature, with the participation and cooperation of partner institutions located in biodiversity-rich countries. For drug-targeted bioprospecting, an industrial partner is a necessity, which will move a discovery into the pipeline toward commercial product.

Central in any mass bioprospecting is the drafting and signing of an international agreement or Memorandum of Understanding (MOU) or MOA, that should cover issues on access to the genetic resources [the biodiversity], on IP related to discovery, on the sharing of the benefits as part of the process (short-term) and in the event of discovery and commercialization of a product (long-term), as well as on the conservation of the biological resources for the future generations. When ethnobotanical or ethnopharmacological approach is utilized, additional specific requirements that relate to prior informed consent (PIC), recognition of Indigenous Intellectual Property (IIP) and Indigenous Intellectual Property Rights (IIPR), as well as short- and long-term benefit-sharing are “priority” items to take into account.

Collection methodology that facilitates recollection, in the event of active species, is essential in order to ensure a reliable supply of biomass for larger scale compound isolation to meet the requirements of in vivo and late-stage preclinical studies. This includes good field documentation, use of Global Positioning System (GPS) to pinpoint site locations, mapping of sites and the ready availability of superior computer database support. As an example, the field data for the UIC ICBG program are posted on the Internet in the “Atlas of Seed Plants of Cuc Phuong National Park” (<http://uic-icbg.pharm.uic.edu>). This information is also made available in hardcopy form (Soejarto et al., 2004c). Good collection

methodology is especially important for marine sites, though collection technology is different (Wright et al., 1996). Needless to say, precise taxonomic identification of organisms involved either in the field, or with subsequent support from taxonomic specialists, is crucial for accurate recollection of organisms or targeted collection of specific organisms (Soejarto, 1996).

Another important tool is plant tissue or cell culture technologies, which are suited for scale-up production of bioactive metabolites once they have been determined to be of interest (DiCosmo and Misawa, 1995; Kirakosyan et al., 2004). In the UIC ICBG, plant cell culture has been applied to eliminate the uncertainty of re-accessing native plant samples that have exhibited interesting chemistry. Further development of sustainable supplies of compounds for clinical trials or commercialized drugs is critical, and may proceed by exploring sustainable harvest methods, cultivation (including aquaculture), microbial fermentation, genetic engineering and semi-synthesis or synthesis of candidate drugs or analogs (Wender et al., 1999). Such effort will ensure adequate supply of the compound while protecting the source organism and its habitat from overexploitation. Planning for sustainability should begin early in product development (Cragg et al., 1993; Cragg, 1998). In the UIC-based Vietnam–Laos ICBG, plant tissue culture is also being used to produce biomass of rare and threatened species at Cuc Phuong National Park, for purposes of both biological evaluation and increasing species population in the park. It is also intended to promote economic development among communities in the ICBG research site by helping members of the community to generate income from distributed starter plants, in cases of economically valuable species.

Acknowledgements

The UIC ICBG bioprospecting project, “Studies on Biodiversity of Vietnam and Laos”, is supported by Grants 1-UO1-TW01015-01 (1998–2003) and 2-UO1-TW001015-06 (2003–2008), International Cooperative Biodiversity Groups Program, at the Fogarty International Center, National Institutes of Health (NIH), through funds from the NIH, National Science Foundation (NSF) and Foreign Agriculture Service of the United States Department of Agriculture (USDA-AFS). The authors express thanks to government authorities of Vietnam and Laos for their cooperation and for the access provided to the biological resources of Vietnam and Laos, respectively, for the ICBG study.

References

Alexiades, M.N. (Ed.), 1996. Selected Guidelines for Ethnobotanical Research: A Field Manual. New York Botanical Garden, Bronx, NY, 306 pp., i–xx.

- Bien, L.K., Hiep, N.T., Loc, P.K., Cuong, N.M., Soejarto, D.D., 2003. The diversity of the flora of Vietnam 14. *Symphorema* Roxb. and *S. involucratum* Roxb., a new genus and species occurrence for the flora of Vietnam. Genetics and Applications 3, 34–38.
- Bien, L.K., Hiep, N.T., Soejarto, D.D., Cuong, N.M., Loc, P.K., 2004. The diversity of the flora of Vietnam 16. Some more data on *Tubocapsicum* (Wettst.) Makino and *T. anomalum* (Franch. et Savat.) Makino (Solanaceae), little known genus and species of the flora. Genetics and Applications 2, 28–32.
- Chien, N.Q., Hung, N.V., Santarsiero, B.D., Mesecar, A.D., Cuong, N.M., Soejarto, D.D., Pezzuto, J.M., Fong, H.H., Tan, G.T., 2004. New 3-O-acyl betulinic acids from *Strychnos vanprukii* Craib. Journal of Natural Products 67, 994–998.
- Cragg, G.M., 1998. Paclitaxel (Taxol): a success story with valuable lessons for natural product drug discovery and development. Medical Research Review 18, 315–331.
- Cragg, G.M., Boyd, M., 1996. Drug discovery and development at the National Cancer Institute: the role of natural products of plant origin. In: Balick, M.J., Elisabetsky, E., Laird, S.A. (Eds.), Medicinal Plant Resources of the Tropical Forest. Columbia University Press, New York, pp. 101–136.
- Cragg, G.M., Schepartz, S.A., Suffness, M., Grever, M.R., 1993. The taxol supply crisis. New NCI policies for handling the large-scale production of novel natural product anticancer and anti-HIV agents. Journal of Natural Products 56, 1657–1668.
- Cragg, G.M., Simon, J.E., Jato, J.G., Snader, K.M., 1996. Drug discovery and development at the National Cancer Institute: potential for new pharmaceutical crops. In: Janick, J. (Ed.), Progress in New Crops. ASHS Press, Arlington, VA, pp. 554–560.
- Dac, L.X., Hai, H.H., Ha, D.T.H., Danh, N.T., Xuan, L.T., Hai, N.V., Binh, L.T., 2004. Micropropagation and biodiversity conservation of *Litsea verticillata* collected from Cuc Phuong National Park by using plant tissue culture technique. Journal of Biotechnology 2, 479–486.
- DiCosmo, F., Misawa, M., 1995. Plant cell and tissue culture: alternatives for metabolite production. Biotechnology Advances 13, 425–453.
- Dietzman, G., Khuong, D.V., Bich, T.Q., Cuong, N.M., Soejarto, D.D., 2002. GIS (Geographic Information System)-based biodiversity assessment as a tool to support conservation initiatives at Cuc Phuong National Park. Advances in Natural Sciences 3, 255–271.
- Dzu, N.V., Hiep, N.T., Loc, P.K., Soejarto, D.D., 2003. Little known species and new records on the occurrence of Angiosperm species in Cuc Phuong National Park III. New records of Aroids (Araceae). Advances in Natural Sciences 4, 1–14.
- Eisner, T., 1989. Prospecting for nature’s chemical riches. Issues in Science and Technology 6, 31–34.
- Eisner, T., 1992. Chemical prospecting: a proposal for action. In: Borhmann, F.H., Kellert, S.R. (Eds.), Ecology, Economics, and Ethics: The Broken Cycle. Yale University Press, New Haven, CT, pp. 196–202.
- Executive Order 247, 1995. Prescribing Guidelines and Establishing a Regulatory Framework for the Prospecting of Biological and Genetic Resources, Their By-Products and Derivatives, for Scientific and Commercial Purposes, and for Other Purposes. Executive Order No. 247, May 18, pp. 1–10 (available at http://www.grain.org/brl_files/philippines-bioprospecting-1995-en.pdf).
- FAO, 1983. Annex to Resolution 8/83. International Undertaking on Plant Genetic Resources. In: Twenty-Second Session of the FAO Conference, Rome (available at <ftp://ext-ftp.fao.org/ag/cgrfa/Res/C8-83E.pdf>).
- Fogarty International Center, 2004. International Cooperative Biodiversity Groups (ICBG) (available at <http://www.fic.nih.gov/programs/icbg.html>).
- Hallock, Y.F., Cragg, G.M., 2003. National Cooperative Drug Discovery Group (NCDDGs): a successful model for public private partnerships in cancer drug discovery. Pharmaceutical Biology 41, 78–91.
- Hammond, P.M., 1995. The record to date: described species. In: Heywood, V.H., Watson, R.T. (Eds.), Global Biodiversity Assessment.

- Cambridge University Press, Cambridge, pp. 113–138 (Chapter 3.1).
- Harshberger, J.W., 1896. The purposes of ethnobotany. *Botanical Gazette* 21, 146–154.
- Hiep, N.T., Loc, P.K., 2000. The diversity of the flora of Vietnam. 4. *Gardneria* Wall. and *G. ovata* Wall., a new genus and species for the flora. *Genetics and Applications* 3, 10–13 (in Vietnamese with an English summary).
- Hiep, N.T., Soejarto, D.D., Loc, P.K., 2002. Tinh da dang cua he thuc vat Viet Nam 10. *Nyctocalos* Teijsm. & Binn. Nuc nac leo (ho Nuc nac Bignoniaceae), chi moi cho he thuc vat Viet Nam. Tap chi Di truyen hoc va ung dung 1, 40–43 (The diversity of the flora of Vietnam 10. *Nyctocalos* Teijsm. & Binn. (family Bignoniaceae), new genus for flora of Vietnam. *Genetics and Applications* 1, 40–43).
- ICBG/AP4, 2005. 50 Medicinal Plants Used by Muong People in Cuc Phuong. Institute of Biotechnology. Associate Program 4, UIC ICBG, Publ., Hanoi (in Vietnamese; postcard-size loose-bound booklet, fully illustrated).
- International Society of Ethnobiology Code of Ethics, 2005. Available at <http://guallart.dac.uga.edu/ISE/iseEthics.html>.
- International Society of Ethnobiology Constitution, 2005. Annex II (available at <http://guallart.dac.uga.edu/ISE/iseConstitution.html>).
- International Society of Ethnopharmacology, 2005. ISE Constitution (available at <http://www.ethnopharmacology.org/iseConst.html>).
- Journal of Ethnopharmacology, 2005. Inside front cover. *Journal of Ethnopharmacology* 98 (inside front cover).
- Kinghorn, A.D., Farnsworth, N.R., Soejarto, D.D., Cordell, G.A., Swanson, S.M., Pezzuto, J.M., Wani, M.C., Wall, M.E., Oberlies, N.H., Kroll, D.J., Kramer, R.A., Rose, W.C., Vite, G.D., Fairchild, C.R., Peterson, R.W., Wild, R., 2003. Novel strategies for the discovery of plant-derived anticancer agents. *Pharmaceutical Biology* 41, 53–67.
- Kirakosyan, A., Sirvent, T.M., Gibson, D.M., Kaufman, P.B., 2004. The production of hypericin and hyperforin by in vitro cultures of St. John's wort (*Hypericum perforatum*). *Biotechnology and Applied Biochemistry* 39, 71–81.
- Loc, P.K., Do, N.T., Hoan, D.T., Hiep, N.T., Cuong, N.M., Soejarto, D.D., 2002. Tinh da dang cua he thuc vat Viet Nam 12. *Phrynium hainanense* T.L. Wu & S.J. Chen La dong che sau (ho Hoang tinh Marantaceae), Loai bo sung cho he thuc vat Viet Nam. Tap chi Di truyen hoc va ung dung 3, 23–26 (The diversity of the flora of Vietnam 12. *Phrynium hainanense* T.L. Wu & S.J. Chen (family Marantaceae), new species for flora of Vietnam. *Genetics and Applications* 3, 23–26).
- Loc, P.K., Hiep, N.T., Soejarto, D.D., Cuong, N.M., 2004. The diversity of the flora of Vietnam 17. *Anodendron howii* Tsiang (Apocynaceae), new species occurrence for the flora. *Genetics and Applications* 3, 36–39.
- Mays, T.D., Mazan, K.D., 1996. Legal issues in sharing the benefits of biodiversity prospecting. *Journal of Ethnopharmacology* 51, 93–109.
- NCI-DTP-GCOB, 2005. NCI-DTP Website at http://dtp.nci.nih.gov/branches/gcob/gcob_web3.html.
- NIH News, 2003. Third Round Awards are Announced Under Interagency Biodiversity Program (available at <http://www.nih.gov/news/pr/dec2003/fic-16.htm>).
- Penso, G., 1976. Inventory of Medicinal Plants and Compilation of a List of The Most Widely Used Plants. Geneva, WHO Document DPM/WP/78.2.
- Reid, W.A., Laird, S.A., Gamez, R., Sittenfeld, A., Janzen, A.H., Gollin, M.A., Juma, C., 1993. A new lease on life. In: Reid, W.A., Laird, S.A., Meyer, C.A., Gamez, R., Sittenfeld, A., Janzen, D.H., Gollin, M.A., Juma, C. (Eds.), *Biodiversity Prospecting: Using Genetic Resources for Sustainable Development*. World Resources Institute, pp. 1–52.
- Riley, M.C., 2001. Traditional Medicine Research Center (TMRC): a potential tool for protecting Traditional and Tribal Medicinal Knowledge in Laos. *Cultural Survival Quarterly* 24, 21–24.
- Rivier, L., Bruhn, J.G., 1979. Editorial. *Journal of Ethnopharmacology* 1, 1.
- Rosenthal, J.P., 1996. Equitable sharing of biodiversity benefits: agreements on genetic resources. In: Presentation at the International Conference on Incentive Measures for the Conservation and the Sustainable Use of Biological Diversity in Cairns, Australia, 25–28 March (The meeting was sponsored and organized by the Organisation for Economic Co-Operation and Development. The paper appears in print form in the publication—Investing in Biological Diversity: Proceedings of the Cairns Conference, OECD 1997, available at <http://www.fic.nih.gov/programs/oecdub.html>).
- Rosenthal, J.P., 1997. Integrating drug discovery, biodiversity conservation, and economic development: early lessons in the International Cooperative Biodiversity Groups. In: Grifo, F., Rosenthal, J. (Eds.), *Biodiversity and Human Health*. Island Press, Washington, DC, pp. 281–301.
- Rosenthal, J.P., Pezzuto, J.M. (Eds.), 1999. *Drug Discovery, Economic Development and Conservation: The International Cooperative Biodiversity Groups*. Swets & Zeitlinger, The Netherlands, pp. 1–144 (*Pharmaceutical Biology* 37, 1–144).
- Rosenthal, J.P., Beck, D.A., Bhat, A., Biswas, J., Brady, L., Bridbord, K., Collins, S., Cragg, G., Edwards, J., Fairfield, A., Gottlieb, M., Gschwind, L.A., Hallock, Y., Hawks, R., Hegyeli, R., Johnson, G., Keusch, G.T., Lyons, E.E., Miller, R., Rodman, J., Roskoski, J., Siegel-Causey, D., 1999. Combining high risk science with ambitious social and economic goals. *Pharmaceutical Biology* 37, 6–21.
- Schippmann, U., Leaman, D.J., Cunningham, A.B., 2002. Impact of cultivation and gathering of medicinal plants on biodiversity: global trends and issues. In: FAO, *Biodiversity and The Ecosystem Approach in Agriculture, Forestry and Fisheries*. Satellite Event on the Occasion of the Ninth Regular Session of the Commission on Genetic Resources for Food and Agriculture. Rome, 12–13 October (Inter-Departmental Working Group on Biological Diversity for Food and Agriculture, Rome, available at http://www.fao.org/documents/show_cdr.asp?url_file=/docrep/005/aa010e/aa010e00.htm).
- Schultes, R.E., von Reis, S. (Eds.), 1995. *Ethnobotany: Evolution of a Discipline*. Chapman & Hall, London, 414 pp.
- Secretariat of the Convention on Biological Diversity, 2001. *Handbook of the Convention on Biological Diversity*. Earthscan Publications Ltd., London/Sterling, VA, pp. 3–26 (available at <http://www.biodiv.org/convention/articles.asp>).
- Soejarto, D.D., 1993. Logistics and politics in plant drug discovery. The other end of the spectrum. In: Kinghorn, A.D., Balandrin, M. (Eds.), *Human Medicinal Agents from Plants*. American Chemical Society, Washington, DC, pp. 98–111.
- Soejarto, D.D., 1996. Biodiversity prospecting and benefit-sharing: perspectives from the field. *Journal of Ethnopharmacology* 51, 1–5.
- Soejarto, D.D., Gyllenhaal, C., Ashton, P.S., Sohmer, S.H., 1996. Plant explorations in Asia under the sponsorship of the National Cancer Institute, 1986–1991: an overview. In: Balick, M.J., Elisabetsky, E., Laird, S.A. (Eds.), *Medicinal Plant Resources of the Tropical Forest*. Columbia University Press, New York, pp. 284–310 (Chapter 21).
- Soejarto, D.D., Gyllenhaal, C., Fong, H.H.S., Xuan, L.T., Hiep, N.T., Hung, N.V., Bich, T.Q., Southavong, S., Sydara, K., Pezzuto, J.M., 2004b. The UIC ICBG (University Of Illinois at Chicago International Cooperative Biodiversity Group) Memorandum of Agreement: a model of benefit-sharing arrangement in natural products drug discovery and development. *Journal of Natural Products* 67, 294–299.
- Soejarto, D.D., Gyllenhaal, C., Regalado, J.C., Pezzuto, J.M., Fong, H.H.S., Tan, G.T., Hiep, N.T., Xuan, L.T., Binh, D.Q., Hung, N.V., Bich, T.Q., Thin, N.N., Loc, P.K., Vu, B.M., Southavong, B.H., Sydara, K., Bouamanivong, S., O'Neill, M.J., Lewis, J., Xie, X.-M., Dietzman, G., 1999. Studies on biodiversity of Vietnam and Laos: the UIC-based ICBG program. *Pharmaceutical Biology* 37, 100–113.
- Soejarto, D.D., Gyllenhaal, C., Regalado, J.C., Pezzuto, J.M., Fong, H.H.S., Tan, G.T., Hiep, N.T., Xuan, L.T., Hung, N.V., Bich, T.Q., Loc, P.K., Vu, B.M., Southavong, B.H., Sydara, K., Bouamanivong, S., O'Neill, M.J., Dietzman, G., 2002a. An international collaborative

- program to discover new drugs from tropical biodiversity of Vietnam and Laos. *Natural Product Science* 8, 1–15.
- Soejarto, D.D., Gyllenhaal, C., Tarzian Sorensen, J.A., Fong, H.H.S., Xuan, L.T., Binh, L.T., Hiep, N.T., Hung, N.V., Vu, B.M., Bich, T.Q., Southavong, B.H., Sydara, K., Pezzuto, J.M., 2004a. Bio-prospecting arrangement: experiences between the north and the south. In: Mahoney, R. (Ed.), *Handbook of Best Practices for Management of Intellectual Property in Health Research and Development*. Centre for the Management of Intellectual Property in Health Research and Development, MIHR, Oxford, UK, pp. 169–190 (third printing).
- Soejarto, D.D., Hiep, N.T., Loc, P.K., Bien, L.K., Dai, T.D., Regalado, J.C., Cuong, N.M., 2002d. Little known species and new records on the occurrence of angiosperm species in Cuc Phuong National Park. 1. *Helicopsis lobata*, *Alangium tonkinense*, *Illigera parviflora*, *Keenania tonkinensis*, *Lysimachia insignis*, and *Kadsura coccinea*. *Advances in Natural Sciences* 3, 57–70.
- Soejarto, D.D., Hiep, N.T., Loc, P.K., Cuong, N.M., Bien, L.K., Dai, T.D., Kadushin, M.R., 2004c. Seed Plants of Cuc Phuong National Park, Vietnam. A Documented Checklist. Cuc Phuong National Park, Ninh Binh, Vietnam. Agricultural Publishing House, Hanoi, Vietnam, pp. i–xxxiv, 1–760 (plates I–XCIX).
- Soejarto, D.D., Regalado, J.C., Dietzman, G., Gyllenhaal, C., Hiep, N.T., Loc, P.K., Huong, N.T., Bich, T.Q., Cuong, N.M., 2001. Development and management of a computerized database of seed plants of Cuc Phuong National park. In: *International Workshop on Biology (Workshop Proceedings)*, vol. 1, Hanoi (Vietnam), 2–5 July, pp. 192–195.
- Soejarto, D.D., Regalado, J.C., Hiep, N.T., Loc, P.K., Bien, L.K., Bich, T.Q., Cuong, N.M., Kadushin, M.R., 2002c. Atlas of Seed Plants of Cuc Phuong National Park (online, posted starting in 2002, available at <http://uic-icbg.pharm.uic.edu>).
- Soejarto, D.D., Xuan, L.T., Vu, B.M., Dac, L.X., Bich, T.Q., Southavong, B.H., Sydara, K., Bouamanivong, S., Zhang, H.J., Fong, H.H.S., Tan, G., Pezzuto, J., Franzblau, S.G., Gyllenhaal, C., Riley, M.C., Hiep, N.T., Loc, P.K., Hung, N.V., 2002b. Implementing IPR and benefit-sharing arrangements: experiences in the University of Illinois at Chicago–Vietnam–Laos ICBG. In: *Intellectual Property Rights and Traditional Knowledge on Genetic Resources in Pharmaceutical and Cosmetic Business*, JBA/NITE, International Symposium 2002, 14 November, Tokyo, Japan, pp. 47–83.
- TMRC, 2003. Bulletin on Lao Traditional Medicine #5. Published by Traditional Medicine Research Center, Ministry of Health, Vientiane, Laos (in Lao language, Bulletin started in 2001).
- Vu, B.M., Xuan, L.T., Hai, N.V., Soejarto, D.D., 2002a. ICBG Associate Program 4 and Buffer Zones of Cuc Phuong National Park. *Vietnam Agriculture* 87, 16 (in Vietnamese).
- Vu, B.M., Xuan, L.T., Hai, N.V., Soejarto, D.D., 2002b. An ICBG/AP4-created model of household economic development for a commune in buffer zone of Cuc Phuong National Park. *Journal of Economic Development Review* 59, 1–3 (in Vietnamese).
- Wender, P.A., Hinkle, K.W., Koehler, M.F., Lippa, B., 1999. The rational design of potential chemotherapeutic agents: synthesis of bryostatin analogues. *Medical Research Review* 19, 388–407.
- Wilson, E.O., Peter, F.M. (Eds.), 1988. *Biodiversity*. National Academy Press, Washington, DC, pp. 3–18.
- Wright, A.D., Konig, G.M., Angerhofer, C.K., Greenidge, P., Linden, A., 1996. Antimalarial activity: the search for marine-derived natural products with selective antimalarial activity. *Journal of Natural Products* 59, 710–716.
- Zhang, H.J., Hung, N.V., Cuong, N.M., Soejarto, D.D., Pezzuto, J.M., Fong, H.H.S., Tan, G.T. Natural anti-HIV constituents from *Litsea verticillata*. *Planta Medica*, in press.
- Zhang, H.-J., Qiu, S.-H., Tamez, P., Tan, G.T., Aydogmus, Z., Hung, N.V., Cuong, N.M., Angerhofer, C., Soejarto, D.D., Pezzuto, J.M., Fong, H.H.S., 2002a. Antimalarial agents from plants II. Decursivine, a new antimalarial indole alkaloid from *Rhaphidophora decursiva*. *Pharmaceutical Biology* 40, 221–224.
- Zhang, H.J., Tamez, P., Aydogmus, Z., Tan, G.T., Saikawa, Y., Hashimoto, K., Nakata, M., Hung, N.V., Xuan, L.T., Cuong, N.M., Soejarto, D.D., Pezzuto, J.M., Fong, H.H.S., 2002b. Antimalarial agents from plants. III. Trichothecenes from *Ficus fistulosa* and *Rhaphidophora decursiva*. *Planta Medica* 68, 1088–1091.
- Zhang, H.-J., Tamez, P.A., Hoang, V.D., Tan, G.T., Hung, N.V., Xuan, L.T., Cuong, N.M., Thao, D.T., Soejarto, D.D., Fong, H.H.S., Pezzuto, J.M., 2001. Antimalarial compounds from *Rhaphidophora decursiva*. *Journal of Natural Products* 64, 772–777.
- Zhang, H.-J., Tan, G.T., Hoang, V.D., Hung, N.V., Cuong, N.M., Soejarto, D.D., Pezzuto, J.M., Fong, H.H.S., 2003. Natural anti-HIV agents. Part III. Litseaverticillols A–H, novel sesquiterpenes from *Litsea verticillata*. *Tetrahedron* 59, 141–148.